

**Intracellular crowding of macromolecules defines the mode and sequence of substrate uptake by *E. coli* and constrains its metabolic activity and growth**

**Supporting Information, Part 3:  
Supplementary Tables**

Q.K. Beg<sup>1\*</sup>, A. Vázquez<sup>2\*</sup>, J. Ernst<sup>3</sup>, M.A. de Menezes<sup>4</sup>, Z. Bar-Joseph<sup>3</sup>, A.-L. Barabási<sup>5</sup> and Z.N. Oltvai<sup>1</sup>

<sup>1</sup>Department of Pathology, University of Pittsburgh, Pittsburgh, PA, 15261, USA

<sup>2</sup>The Simons Center for Systems Biology, Institute for Advanced Study, Princeton, NJ 08540, USA

<sup>3</sup>Machine Learning Department, Carnegie-Mellon University, Pittsburgh, PA, 15217, USA

<sup>4</sup>Instituto de Física, Universidade Federal Fluminense, Rio de Janeiro, 24210, Brazil.

<sup>5</sup>Department of Physics and Center for Complex Networks Research, University of Notre Dame, Notre Dame, IN 46556, USA

**CONTENT**

**Table S1 (separate excel table-Supporting file 4):** List of calculated  $a_i$  for several *E. coli* enzymes

**Table S2:** Full names of genes, enzymes their functions and substrates shown in Figure 3

**Table S3: (separate excel table-Supporting file 5):** Time series microarray data of cultures shown in Figure 2, 3 and 4

**Table S4a: (separate excel table-Supporting file 6):** Microarray data of single carbon-limited batch-cultures, described in Figure 4 (bottom panel)

**Table S4b: (separate excel table-Supporting file 7):** Microarray data of glucose and lactate-limited batch-cultures, described in Figure 4 (bottom panel)

## Supplementary Tables

**Table S2:** Major functions and reactions catalyzed by the transporters/enzymes encoded by the genes involved in substrate uptake, as shown in Figure 3 of the manuscript

Gene and carbon source	Functions and reactions catalyzed by enzymes/proteins
<b>Glucose</b>	
<i>ptsG</i>	<p><u>Functions:</u> The product of <i>ptsG</i> gene (glucose-specific PTS permease) is responsible for uptake of exogenous glucose from the medium, releasing the phosphate ester into the cell cytoplasm in preparation for metabolism, primarily via glycolysis (Buhr and Erni 1993; Postma et al. 1993). PtsG/Crr, the glucose-specific PTS permease, belongs to the functional superfamily of the phosphoenolpyruvate (PEP)-dependent, sugar transporting phosphotransferase system (PTS). The PTS transports and simultaneously phosphorylates its sugar substrates in a process called group translocation.</p> <p><u>Reaction:</u> phosphoenolpyruvate + <math>\beta</math>-D-glucose[periplasmic space] <math>\rightarrow</math> <math>\beta</math>-D-glucose-6-phosphate + pyruvate</p>
<b>Maltose</b>	
<i>malEFGK</i>	<p><u>Functions:</u> <i>malKFGK</i> operon plays the major role in the maltose transport system and it belongs to the ATP-Binding Cassette (ABC) superfamily of transporters (Wu and Mandrand-Berthelot 1995). <i>malE</i> is the periplasmic maltose-binding protein, <i>malF</i> and <i>malG</i> are the integral membrane components of the ABC transporter, and <i>malK</i> is the ATP-binding component of the ABC transporter (MalFGK<sub>2</sub>) (Boos and Shuman 1998; Joly et al. 2004)</p> <p><u>Reactions:</u> maltose<sub>[extracellular space]</sub> <math>\Leftrightarrow</math> maltose<sub>[cytosol]</sub> ATP + maltose<sub>[periplasmic space]</sub> + H<sub>2</sub>O <math>\Leftrightarrow</math> ADP + phosphate + maltose<sub>[cytosol]</sub></p>
<i>malQ</i>	<p><u>Functions:</u> <i>malQ</i> codes for amylomaltase, which is responsible for degrading maltose after transport into the cell (Neidhardt et al. 1996). The glucose liberated in the degradation reaction is then used in glycolysis. Amylomaltase also recognizes maltotriose and larger maltodextrins (donors), cleaving off the reducing glucose residue and transferring the remaining dextrinyl residue onto the nonreducing end of maltodextrin (acceptors), including maltose and glucose. Amylomaltase thus produces glucose and longer maltodextrins from maltotriose, the smallest donor substrate, as well as from longer linear maltodextrins (Boos and Shuman 1998)</p> <p><u>Reaction:</u></p>

	$\text{H}_2\text{O} + \text{maltose} \rightleftharpoons 2 \beta\text{-D-glucose}$ $\text{maltotriose} + \text{maltose} \rightleftharpoons \text{maltotetraose} + \beta\text{-D-glucose}$
<i>glk</i>	<u>Functions:</u> Under normal conditions, glucokinase plays a minor role in <i>E. coli</i> glucose metabolism. Under anabolic stress the enzyme is required to supplement the levels of glucose 6-phosphate. <u>Reactions:</u> $\beta\text{-D-glucose} + \text{ATP} \rightleftharpoons \beta\text{-D-glucose-6-phosphate} + \text{ADP}$
<b>Galactose</b>	
<i>mglABC</i>	<u>Functions:</u> MglABC is a beta-methylgalactoside transport system that is a member of the ATP-Binding Cassette (ABC) superfamily of transporters (Wu and Mandrand-Berthelot 1995). The <i>mglB</i> gene codes for a galactose-binding protein that serves both as the galactose chemoreceptor as well as the recognition component of the $\beta$ -methylgalactoside transport system, which utilizes the galactose-binding protein; <i>mglC</i> encodes the integral membrane component; and <i>mglA</i> encodes the ATP-binding component of the ABC transporter (Harayama et al. 1983). <u>Reaction:</u> $\text{ATP} + \beta\text{-D-galactose}_{[\text{periplasmic space}]} + \text{H}_2\text{O} \rightleftharpoons \text{ADP} + \text{phosphate} + \beta\text{-D-galactose}_{[\text{cytosol}]}$
<i>galE</i>	<u>Functions:</u> <i>galE</i> codes for UDP-galactose 4-epimerase, which catalyzes a hydride transfer and the interconversion of UDP-galactose and UDP-glucose as part of galactose catabolism (Swanson and Frey 1993). <u>Reaction:</u> $\text{UDP-D-glucose} \rightleftharpoons \text{UDP-galactose}$
<i>galK</i>	<u>Functions:</u> Galactokinase, coded by <i>galK</i> catalyzes the first step in the galactose metabolism <u>Reaction:</u> $\text{D-galactose} + \text{ATP} = \alpha\text{-D-galactose 1-phosphate} + \text{ADP}$
<i>galT</i>	<u>Functions:</u> <i>galT</i> codes for galactose 1-phosphate uridylyltransferase, which catalyzes an interconversion reaction in galactose catabolism <u>Reaction:</u> $\text{UDP-D-glucose} + \alpha\text{-D-galactose 1-phosphate} = \alpha\text{-D-glucose 1-phosphate} + \text{UDP-galactose}$ $\alpha\text{-D-galactose 1-phosphate} + \text{UTP} = \text{UDP-galactose} + \text{diphosphate}$
<i>galP</i>	<u>Functions:</u>

	<p>GalP is one of two, along with MglABC, major routes for galactose transport into <i>E. coli</i>. 2-deoxy- D-galactose is a specific substrate for GalP but not for MglABC and GalP operates by a sugar-proton symport mechanism while MglABC does not.</p> <p><u>Reaction:</u>  <math>H^+_{[periplasmic\ space]} + \beta\text{-D-galactose}_{[periplasmic\ space]} \Leftrightarrow H^+_{[cytosol]} + \beta\text{-D-galactose}_{[cytosol]}</math></p>
<i>pgm</i>	<p><u>Functions:</u>  <i>pgm</i> codes for phosphoglucose mutase, which catalyzes conversion of glucose-1-phosphate to glucose-6-phosphate. Maximum activity is only obtained in the presence of <math>\alpha</math>-D-glucose 1,6-bisphosphate. This bisphosphate is an intermediate in the reaction, being formed by transfer of a phosphate residue from the enzyme to the substrate, but the dissociation of bisphosphate from the enzyme complex is much slower than the overall isomerization.</p> <p><u>Reaction:</u>  <math>\alpha\text{-D-glucose-1-phosphate} \rightarrow \alpha\text{-D-glucose-6-phosphate}</math></p>
<b>Glycerol</b>	
<i>glpK</i>	<p><u>Function:</u>  <i>glpK</i> codes for glycerol kinase, which catalyzes the MgATP-dependent phosphorylation of glycerol to yield sn-glycerol 3-phosphate (Pattigrew et al. 1988). This is also the rate limiting step in glycerol utilization in <i>E. coli</i>.</p> <p><u>Reaction:</u>  <math>\text{glycerol} + \text{ATP} \rightarrow \text{sn-glycerol-3-phosphate} + \text{ADP}</math></p>
<i>glpF</i>	<p><u>Function:</u>  The glycerol facilitator, GlpF, allows the facilitated diffusion of glycerol into the cell (Stroud et al. 2003).</p> <p><u>Reaction:</u>  <math>\text{glycerol}_{[periplasmic\ space]} \Leftrightarrow \text{glycerol}_{[cytosol]}</math></p>
<i>gpsA</i>	<p><u>Functions:</u>  <i>gpsA</i> codes for glycerol-3-phosphate-dehydrogenase-[NAD(P)<sup>+</sup>], which catalyzes the NAD(P)H-dependent reduction of the glycolytic intermediate dihydroxyacetone-phosphate to produce glycerol-3-phosphate (Lin 1976)</p> <p><u>Reaction:</u>  <math>\text{sn-glycerol-3-phosphate} + \text{NAD(P)}^+ = \text{dihydroxyacetone phosphate} + \text{NAD(P)H} + \text{H}^+</math></p>
<b>Lactate</b>	
<i>lldP</i>	<p><u>Function:</u>  LldP (or LctP) is a lactate/proton symporter responsible for the uptake of L-lactate. The <i>lldP/lctP</i> gene is located in a lactate-inducible operon with the <i>lctD</i> and <i>lctR</i> genes encoding a lactate dehydrogenase and a regulatory protein, respectively (Dong et al. 1993).</p> <p><u>Reaction:</u>  <math>H^+_{[periplasmic\ space]} + \text{lactate}_{[periplasmic\ space]} \Leftrightarrow H^+_{[cytosol]} + \text{lactate}_{[cytosol]}</math></p>

<i>dld</i>	<p><u>Function:</u>  <i>dld</i> codes for D-lactate dehydrogenase. There are three lactate dehydrogenase enzymes in <i>E. coli</i> which interconvert pyruvate and lactate. One is an NAD-linked fermentative dehydrogenase. The other two are membrane-bound flavoproteins, each specific for the D- or L-isomer, and involved in the aerobic respiratory chain of <i>E. coli</i>. The D-lactate dehydrogenase is coded for by the <i>dld</i> gene and it is the primary source of energy to drive the active transport of certain sugars and amino acids into the cell (Matsushita et al. 1986).</p> <p><u>Reaction:</u>  ubiquinone-8 + D-lactate <math>\Leftrightarrow</math> ubiquinol-8 + pyruvate</p>
<b>Acetate</b>	
<i>ackA</i>	<p><u>Function:</u>  The <i>ackA</i> gene product has propionate kinase activity as well as acetate kinase activity. It is unclear whether the two <i>ack</i> genes, <i>ackA</i> and <i>ackB</i>, code for two distinct acetate kinase enzymes or control a single enzyme. Helps in conversion of acetate to acetyl phosphate. The <i>ackA</i> encoded propionate kinase 2 has an important role in propionyl-CoA metabolism (Hesslinger et al. 1998). Acetate kinase can also catalyze acetylation of CheY, increasing signal strength for flagellar rotation. (Barak et al. 1998)</p> <p><u>Reactions:</u>  ATP + propionate <math>\Leftrightarrow</math> ADP + propionyl-P  acetate + ATP <math>\Leftrightarrow</math> acetylphosphate + ADP</p>
<i>pta</i>	<p><u>Function:</u>  <i>pta</i> gene codes for phosphate acetyltransferase, which can utilize both acetyl-CoA and propionyl-CoA.</p> <p><u>Reactions:</u>  phosphate + acetyl-CoA <math>\Leftrightarrow</math> acetylphosphate + coenzyme A  propionyl-CoA + phosphate <math>\Leftrightarrow</math> propionyl-P + coenzyme A</p>
<i>acs</i>	<p><u>Function:</u>  <i>acs</i> gene codes for acetyl-CoA synthetase. There are two distinct pathways by which <i>E. coli</i> converts acetate to acetyl-CoA. Acetyl-CoA synthetase catalyzes one of them. It is thought that this ACS pathway functions in a mainly anabolic role, scavenging acetate present in the extracellular medium (Kumari et al. 1995). ACS also can catalyze acetylation of CheY, increasing signal strength for flagellar rotation (Ramakrishna et al. 1998).</p> <p><u>Reactions:</u>  coenzyme A + 4-coumarate + ATP <math>\rightarrow</math> coumaroyl-CoA + diphosphate + AMP  coenzyme A + propionate + ATP <math>\Leftrightarrow</math> propionyl-CoA + diphosphate + AMP  coenzyme A + acetate + ATP <math>\Leftrightarrow</math> acetyl-CoA + diphosphate + AMP</p>

## References

1. Barak R, Abouhamad WN, Eisenbach M (1998). Both acetate kinase and acetyl coenzyme A synthetase are involved in acetate-stimulated change in the direction of flagellar rotation in *Escherichia coli*. J Bacteriol 180; 985-988
2. Boos W, Shuman H (1998). Maltose/maltodextrin system of *Escherichia coli*: transport, metabolism, and regulation. Microbiol Mol Biol Rev 62; 204-229
3. Buhr A, Erni B (1993). Membrane topology of the glucose transporter of *Escherichia coli*. J Biol Chem 268; 11599-11603
4. Dong JM, Taylor JS, Latour DJ, Iuchi S, Lin EC (1993). Three overlapping *lct* genes involved in L-lactate utilization by *Escherichia coli*. J Bacteriol 175; 6671-6678
5. Harayama S, Bollinger J, Iino T, Hazelbauer GL (1983). Characterization of the *mgl* operon of *Escherichia coli* by transposon mutagenesis and molecular cloning. J Bacteriol 153; 408-415
6. Hesslinger C, Fairhurst SA, Sawers G (1998). Novel keto acid formate-lyase and propionate kinase enzymes are components of an anaerobic pathway in *Escherichia coli* that degrades L-threonine to propionate. Mol Microbiol 27; 477-492
7. Joly N, Bohm A, Boos W, Richet E (2004). MalK, the ATP-binding cassette component of the *Escherichia coli* maltodextrin transporter, inhibits the transcriptional activator MalT by antagonizing inducer binding. J Biol Chem 279; 33123-33130
8. Kumari S, Tishel R, Eisenbach M, Wolfe AJ (1995). Cloning, characterization, and functional expression of *acs*, the gene which encodes acetyl coenzyme A synthetase in *Escherichia coli*. J Bacteriol 177; 2878-2886
9. Lin EC (1976). Glycerol dissimilation and its regulation in bacteria. Annu Rev Microbiol 30; 535-578
10. Matsushita K, Kaback HR (1986). D-lactate oxidation and generation of the proton electrochemical gradient in membrane vesicles from *Escherichia coli* GR19N and in proteoliposomes reconstituted with purified D-lactate dehydrogenase and cytochrome o oxidase. Biochemistry 25; 2321-2327
11. Neidhardt FC, Curtiss III R, Ingraham JL, Lin ECC, Low Jr KB, Magasanik B, Reznikoff WS, Riley M, Schaechter M, Umberger HE " *Escherichia coli* and *Salmonella*, Cellular and Molecular Biology, 2<sup>nd</sup> Edition. American Society for Microbiology, Washington, D.C., 1996.
12. Pettigrew DW, Ma DP, Conrad CA, Johnson JR (1988). *Escherichia coli* glycerol kinase. Cloning and sequencing of the *glpK* gene and the primary structure of the enzyme. J Biol Chem 263; 135-139

13. Postma PW, Lengeler JW, Jacobson GR (1993). Phosphoenolpyruvate:carbohydrate phosphotransferase systems of bacteria. *Microbiol Rev* 57; 543-594
14. Ramakrishnan R, Schuster M, Bourret RB (1998). Acetylation at Lys-92 enhances signaling by the chemotaxis response regulator protein CheY. *Proc Natl Acad Sci USA* 95; 4918-4923
15. Stroud RM, Miercke LJ, O'Connell J, Khademi S, Lee JK, Remis J, Harries W, Robles Y, Akhavan D (2003). Glycerol facilitator GlpF and the associated aquaporin family of channels. *Curr Opin Struct Biol* 13; 424-431
16. Swanson BA, Frey PA (1993). Identification of lysine 153 as a functionally important residue in UDP-galactose 4-epimerase from *Escherichia coli*. *Biochemistry* 32; 13231-13236.
17. Wu LF, Mandrand-Berthelot MA (1995). A family of homologous substrate-binding proteins with a broad range of substrate specificity and dissimilar biological functions. *Biochimie* 77; 744-750